

Paravertebral Block to Reduce the Incidence of New Onset Atrial Fibrillation After Cardiac Surgery: A Prospective Randomized Controlled Pilot Trial

Status: Recruiting

Eligibility Criteria

Age: 18 years and over

This study is NOT accepting healthy

Healthy Volunteers: volunteers

Inclusion Criteria:

- undergoing one of the following elective or urgent (but not emergency) surgeries: A) Primary Coronary Artery Bypass Graft (CABG) B) Primary Surgical Aortic Valve Replacement (sAVR) C) Primary Surgical Mitral Valve Replacement (sMVR) D) Combined CABG & surgical valve replacement

Exclusion Criteria:

- history of atrial fibrillation or flutter - Infective endocarditis - Left ventricular ejection fraction (LVEF) < 30% - redo surgery - unable to have a block because of local anesthetic allergy, bleeding problem - Body mass index > 35kg/m² - woman who is pregnant

Conditions & Interventions

Conditions:

Heart & Vascular

Keywords:

Clinics and Surgery Center (CSC), AF, Atrial Fibrillation, CABG, Cardiac Disease, Coronary Artery By-Pass Surgery

More Information

Description: To determine if a perioperative infusion of 0.2% ropivacaine via bilateral T3 paravertebral catheters can decrease the incidence of new onset atrial fibrillation following primary CABG and/or valve surgery and compare a number of secondary outcomes.

Contact(s): Candace Nelson - nelso377@umn.edu

Principal Investigator: James Flaherty

IRB

Number: STUDY00009938

System ID: 29879

Thank you for choosing StudyFinder. Please visit <http://studyfinderstaging.umn.edu> to find a Study which is right for you and contact sfinder@umn.edu if you have questions or need assistance.